



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

blade, while its terminal limit is difficult to determine, because growth gradually decreases toward the tip of the blade.

A second paper, by Miss SHELDON,⁹ deals with the growth of the stipe. It was found that the region of maximum growth in mature plants is 2-4 feet below the blade, the rate decreasing in both directions. In July the rate of growth was found to be about an inch a day.—J. M. C.

Ferns of the oriental tropics.—The richness of the fern vegetation of the tropics is being emphasized by the work of COPELAND.¹⁰ In the papers cited, new species are described in the following genera: *Adiantum*, *Aglaomorpha* (a new subgenus *Holostachyum*, including two other species, being described), *Angiopteris*, *Asplenium*, *Athyrium* (6), *Balantium*, *Cyathea* (6), *Davallia*, *Dryopteris* (6), *Leptochilus*, *Loxogramme* (2), *Marattia*, *Microlepia*, *Oleandra*, *Ophioglossum*, *Polypodium* (8), *Prosaptia*, *Pteris*, *Tectaria* (4), *Trichomanes*, and *Vittaria*. The genera *Diplora* and *Triphlebia* were found to be invalid, because founded upon inconstant and "illusory" features, and should be included in *Phyllitis*, which comprises only three well defined species in the Malay-Polynesian region.—J. M. C.

Sphagnum bogs of Alaska.—RIGG¹¹ has noted the peculiarities of the flora of some Alaskan peat bogs and finds that while sphagnum occurs in many different habitats in Alaska, only where there is an absence of drainage do bogs accompany it. The peat in the bogs visited had a maximum depth of only 2.5 ft. Aside from the sphagnum, *Empetrum nigrum* is the most abundant and uniform in its occurrence, but *Ledum palustre*, *Kalmia glauca*, *Oxycoccus oxycoccus*, and *Drosea rotundifolia* are among other characteristic species. The bogs occur surrounded by treeless areas, by tundras, or by coniferous forests, and vary much in area.—GEO. D. FULLER.

⁹ SHELDON, SARAH M., Notes on the growth of the stipe of *Nereocystis Luetkeana*. *Ibid.* 1:15-18. 1915.

¹⁰ COPELAND, E. B., The ferns of the Batu Lawi Expedition. Appendix III. Roy. Agric. Soc. (Straits Branch). no. 63. 71-72. 1912.

———, Some ferns of northeastern Mindanao. *Leaflets Philipp. Bot.* 5:1679-1684. 1913.

———, Notes on some Javan ferns. *Philipp. Jour. Sci.* 8:139-145. *pls.* 2-4. 1913.

———, On *Phyllitis* in Malaya and the supposed genera *Diplora* and *Triphlebia*. *Ibid.* 8:147-155. *pls.* 5-7. 1913.

———, New Papuan ferns. *Ibid.* 9:1-9. 1914.

———, New Sumatran ferns. *Ibid.* 9:227-233. 1914.

¹¹ RIGG, G. B., Notes on the flora of some Alaskan sphagnum bogs. *Plant World* 17:176-183. 1914.